



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE VOLUME OF SOME SOLIDS.

From *Journal de Mathematiques*.

TRANSLATED BY WILLIAM HOOVER, A. M., DAYTON, OHIO.

A BODY is comprised between two parallel planes, and a section made in this body by a plane parallel to the bases, at a distance x from the lower base, has an area equal to $a + \beta x + \gamma x^2 + \omega x^3$, (1)

a, β, γ, ω being any four constant values, that is to say independent of x .

To determine the volume of this body.

Cut the body by an infinity of equidistant planes parallel to the given planes. In designating by $S_0, S_1, S_2 \dots S_n$ the sections formed, S_0 and S_n designating the planes of the bases, the altitude h would be divided by these planes into n equal parts.

By virtue of the principle of Cavalieri (any body comprised between two parallel planes can be considered as the sum of an infinity of prisms or of cylinders of infinitely small altitude) the volume of the body considered would be expressed by

$$V = \frac{h}{n} (S_0 + S_1 + S_2 \dots + S_{n-1}), \text{ or } V = \frac{h}{n} \Sigma S. \quad (2)$$

This fixed, evaluate the sections $S_0, S_1, S_2 \dots$. If $x = 0, S_0 = a$, $x = \frac{h}{n}, S_1 = a + \beta \frac{h}{n} + \gamma \frac{h^2}{n^2} + \omega \frac{h^3}{n^3}$, $x = \frac{2h}{n}, S_2 = a + \beta \frac{2h}{n} + \gamma \frac{4h^2}{n^2} + \omega \frac{8h^3}{n^3}$,

$$x = \frac{3h}{n}, S_3 = a + \beta \frac{3h}{n} + \gamma \frac{9h^2}{n^2} + \omega \frac{27h^3}{n^3},$$

$$- \quad - \quad - \quad - \quad - \quad - \quad - \quad - \quad - \quad - \\ x = \frac{(n-1)h}{n}, S_{n-1} = a + \beta \frac{(n-1)h}{n} + \gamma \frac{(n-1)^2 h^2}{n^2} + \omega \frac{(n-1)^3 h^3}{n^3};$$

thence $\Sigma S = na + \beta \frac{h}{n} [1 + 2 + 3 \dots + (n-1)] + \gamma \frac{h^2}{n^2} [1^2 + 2^2 + 3^2 \dots + (n-1)^2]$

$$+ \omega \frac{h^3}{n^3} [1^3 + 2^3 + 3^3 \dots + (n-1)^3], = na + \beta \frac{h}{n} \left[\frac{n(n+1)}{2} \right] \\ + \gamma \frac{h^2}{n^2} \left[\frac{n(n+1)}{6} \frac{(2n+1)}{3} \right] + \omega \frac{h^3}{n^3} \left[\frac{n^2(n+1)^2}{4} \right].$$

Replacing in (2) ΣS by this value, we have

$$V = ah + \beta \frac{h^2}{2} \left(1 - \frac{1}{n} \right) + \gamma \frac{h^3}{6} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \omega \frac{h^4}{4} \left(1 - \frac{1}{n} \right)^2.$$

When n increases indefinitely, we have

$$V = ah + \beta \frac{1}{2} h^2 + \gamma \frac{1}{6} h^3 + \omega \frac{1}{4} h^4. \quad (A)$$

REMARK.—If we designate by B , B_1 the upper and lower bases of a body, and by B_2 the section equidistant from the bases we have

$$x = 0, B = a, x = h, B_1 = a + \beta h + \gamma h^2 + \omega h^3,$$

$$x = \frac{1}{2}h, B_2 = a + \beta \frac{1}{2}h + \gamma \frac{1}{4}h^2 + \omega \frac{1}{8}h^3; \text{ whence}$$

$$4B_2 = 4a + 2\beta h + \gamma h^2 + \frac{1}{2}\omega h^3, \text{ and consequently}$$

$$B + B_1 + 4B_2 = 6a + 3\beta h + 2\gamma h^2 + \frac{3}{2}\omega h^3,$$

which multiplied by $\frac{1}{6}h$ produces formula (A). We then have

$$V = \frac{1}{6}h(B + B_1 + 4B_2), \text{ a known formula.}$$

For the Sphere.—If in (1) we make $a = 0$, $\beta = 2\pi R$, $\gamma = -\pi$, $\omega = 0$, that function reduces to

$$2\pi Rx - \pi x^2 = \pi x(2R - x),$$

which represents the area of a circle made in a sphere, at the altitude x .

In this case $B = 0$, $B_1 = 0$, $B_2 = \pi R^2$, $h = 2R$; thence

$$V = \frac{1}{3}R \cdot 4\pi R^2 = \frac{4}{3}\pi R^3.$$

For the spherical segment.—Making $B = 0$, $B_1 = \pi h(2R - h)$, $B_2 = \pi \cdot \frac{1}{2}h(2R - \frac{1}{2}h)$, we have

$$V = \pi \cdot \frac{1}{6}h^2(2R - h + 4R - h) = \frac{1}{3}\pi h^2(3R - h).$$

For the cone.—If h is the altitude of a cone, R the radius of the base, the radius made at a height x is $R' = R - (R \div h)x$; whence

$$\pi R'^2 = \pi R^2 - \frac{2\pi R^2}{h}x + \frac{\pi R^2}{h^2}x^2,$$

the expression to which (1) reduces itself when $a = \pi R^2$, $\beta = -2\pi R^2 \div h$, $\gamma = \pi R^2 \div h^2$, $\omega = 0$. Thence the cone $= \frac{1}{6}h(\pi R^2 + \pi R^2) = \frac{1}{3}\pi R^2 h$.

In the same way we obtain for the volume of the frustum of a cone

$$V = \frac{1}{6}h[\pi R^2 + \pi R^2 + \pi(r + R)^2] = \frac{1}{3}\pi h(R^2 + r^2 + Rr).$$

NOTE ON BILINEAR TANGENTIAL COORDINATES.

BY PROF. F. H. LOUD, COLORADO SPRINGS, COLORADO.

TANGENTIAL equations are usually written as homogeneous, and so analogous to trilinear equations; but interesting relations to the ordinary Cartesian coordinates are thus unnoticed. If the general equation of the first degree in two variables be divided through by its absolute term, and signs changed, it takes the form

$$px + qy - 1 = 0.$$

In the Cartesian system the two coefficients of this equation, p and q , are the reciprocals of the intercepts of its locus upon the two axes; and a recip-